

QUALITY AND STABILITY OF CANOLA AND SUNFLOWER OILS BLENDED WITH PALM OLEIN DURING FRYING AND MICROWAVE HEATING

ZAHRAU BAMALLI NOURUDDEEN

A thesis submitted in fulfilment of the
requirements for the award of the degree of
Master of Engineering (Bioprocess)

Faculty of Chemical Engineering
Universiti Teknologi Malaysia

JANUARY 2013

To

My beloved parents

Muhammad Nourah Bamalli and Sa'adatu Bamalli

My dearest husband

Dr. Nouruddeen Bashir Umar

For your unwavering love, sacrifices, encouragement and best wishes

ACKNOWLEDGEMENT

First and foremost my gratitude goes to my supervisors Dr. Md. Abbas Ali and Assoc. Prof. Dr. Ida Idayu Muhamad for the humble supervision and academic guidance I received from them in the course of my master's programme. May the sky be your limits in all your future endeavours.

My gratitude also goes to Mr. Razam Abdul Latip and Ms. Noor Hidayu Othman for their immense assistance during my laboratory work at Sime Darby R & D centres, Banting and Carey Island, Selangor.

This acknowledgment will be incomplete if I do not mention the unconditional support and well wishes I have received from my in-laws Dr. Bashir Umar and Umma Bashir, my son, Abdulhaleem, my siblings, Halimah, Munira, Amina, Mahmoud and Al-Mustpha, my brothers- and sisters-in law, Salim, Maryam, Ahmad, Hafsa, Munir and Umar. Your encouragement really gave me hope.

Last but not least, I also would like to acknowledge the fun and happy moments I had with FOBERG (Food and Biomaterial Engineering Research Group) postgraduate students. Those moments have really lessened my homesickness.

ABSTRACT

The aim of this study was to evaluate the changes in quality and oxidative degradation of canola oil (CO) and sunflower oil (SFO) and their blends with comparatively low priced palm olein (PO) during frying and microwave heating. The oil blends were prepared by blending PO with CO and SFO in the volume ratios of 20:80 (PC1 and PSF1) and 40:60 (PC2 and PSF2). Refractive indices of the samples increased insignificantly as the heating time performed by the oils increased. The amounts of free fatty acids were found to be higher in PO and in blends compared to the substrate oils. Peroxide values during frying or microwave heating of all the samples were increased until a maximum is reached and then decreased until the end of the heating. Other quality indices such as *p*-anisidine value, polymer content, viscosity, specific extinctions, food oil sensor and color unit values increased significantly in most of the samples as heating time progressed with increase in the blends being lower than those of the substrate oils. The amount of total polar compound in all the oil samples increased significantly with the time increment. During frying and heating, progressive decreases in iodine value and oxidative stability index were detected. However, PO addition to CO or SFO decelerated the formation of primary and secondary oxidative products and polymeric compounds during heating. In Gas Liquid Chromatography (GLC) analysis, the percentage of linoleic and linolenic acids tended to decrease, whereas the percentage of palmitic, stearic and oleic acids increased in most of the samples during frying and microwave heating. The decreased amounts in $C_{18:2}/C_{16:0}$ ratios from the initial values were observed for PO, CO and SFO in frying and microwave heating; being higher than those of their blends. In differential scanning calorimetry (DSC) analysis, addition of PO altered the shapes of endo- or exotherm peaks of CO and SFO. In this study, most of the degradation indicators suggested that the degradation was the fastest in CO or SFO as compared to their blends. Overall, it can be concluded that addition of PO to CO or SFO slowed down the oxidative degradation rate of the substrate oils. Blend PC2 and PSF2 showed slightly better frying performance compared to PC1 and PSF1.

ABSTRAK

Tujuan kajian ini adalah untuk mengkaji perubahan dari segi kualiti dan degradasi oksidatif minyak canola (CO) dan minyak bunga matahari (SFO) dan percampurannya dengan olein minyak sawit yang berharga murah (PO) semasa menggoreng dan di panaskan dalam ketuhar gelombang mikro. Minyak ini di hasilkan dengan mencampurkan PO dengan CO dan SFO dalam nisbah isipadu 20:80 (PC1 dan PSF1) dan, 40:60 (PC2 dan PSF2). Indeks biasan daripada sampel menunjukkan peningkatan ketara apabila suhu pemanasan minyak meningkat. Jumlah asid lemak didapati tinggi didalam PO dan campurannya jika di bandingkan dengan minyak substrat. Nilai peroksida semasa menggoreng atau pemanasan dalam gelombang mikro untuk kesemua sampel meningkat sehingga nilai maksimum di capai dan kemudian menurun sehingga ke akhir pemanasan. Indeks kualiti lain seperti nilai *p*-Anisidina, kandungan polimer, kelikatan, keluputan spesifik, nilai sensor minyak makanan dan unit berwarna bertambah dengan ketara dalam kebanyakan sampel apabila suhu pemanasan progresif meningkat dengan peningkatan campuran yang lebih rendah daripada minyak substrat. Jumlah sebatian terketutub (TPC) dalam semua sampel minyak meningkat dengan ketara dengan peningkatan masa. Semasa menggoreng dan pemanasan, penurunan progresif bagi nilai iodin dan indeks kestabilan oksidatif telah dikesan. Walaubagaimanapun, pertambahan PO kepada CO atau SFO melambatkan pembentukan produk oksidatif utama dan sekunder dan gabungan polimerik semasa pemanasan. Dari analisa gas cecair kromatografi (GLC), peratus asid linoleik dan linolenik semakin menurun, manakala peratus asid palmitik, stearik dan oleik bertambah dalam kebanyakan sampel semasa menggoreng dan pemanasan gelombang mikro. Jumlah nisbah $C_{18:2}/C_{16:0}$ berkurang dari nilai permulaan, masing masing didapati untuk PO, CO dan SFO dalam gorengan dan pemanasan gelombang mikro; didapati lebih tinggi daripada campurannya. Dalam pengimbasan pembezaan kalorimetri (DSC), tambahan PO mengubah bentuk puncak endo dan eksoterma CO dan SFO. Dalam kajian ini, kebanyakan penunjuk degradasi mencadangkan degradasi terpantas adalah dalam CO atau SFO jika di bandingkan dengan campurannya. Secara keseluruhan kajian ini menunjukkan bahawa pertambahan PO kepada CO atau SFO melambatkan kadar degradasi oksidatif oleh minyak substrat. Campuran PC2 dan PSF2 menunjukkan sedikit kemajuan minyak goreng berbanding PC1 dan PSF1.

TABLE OF CONTENTS

CHAPTER	TITLE	PAGE
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENT	iv
	ABSTRACT	v
	ABSTRAK	vi
	TABLE OF CONTENTS	vii
	LIST OF TABLES	xii
	LIST OF FIGURES	xiv
	LIST OF ABBREVIATIONS AND SYMBOLS	xvi
	LIST OF APPENDICES	xviii
1	INTRODUCTION	1
	1.1 Background	1
	1.2 Problem Statement	3
	1.3 Objective of Study	4
	1.4 Scope of Study	5
	1.5 Significance of Study	5
	1.6 Work Flow Chart	6
	1.7 Thesis Outlines	7
2	LITERATURE REVIEW	8
	2.1 Introduction	8
	2.2 A brief description on Palm Olein, Canala Oil and Sunflower Oil	9
	2.2.1 Palm Olein	9

2.2.1.1	Health Benefits of Palm Olein	12
2.2.2	Canola Oil	13
2.2.2.1	Health Benefits of Canola Oil	15
2.2.3	Sunflower Oil	15
2.2.3.1	Health Benefits of Sunflower Oil	18
2.3	Deep-fat Frying	19
2.3.1	Frying Oils	20
2.3.2	Frying Process	21
2.3.3	Heat Transfer during Frying	22
2.3.4	Chemical Reactions during Frying/Heating	24
2.3.4.1	Hydrolysis of Oil	24
2.3.4.2	Oxidations of Oil	25
2.3.4.3	Polymerization of Oil	27
2.3.4.4	Colour Formation	28
2.3.5	Factors Affecting the Quality of Oil during Deep- Fat Frying	29
2.3.5.1	Types of Fryer	29
2.3.5.2	Fat Turnover	28
2.3.5.3	Frying Time and Temperature	30
2.3.5.4	Quality of Frying Oil	30
2.3.5.5	Flavour Quality of Frying Oil	31
2.3.5.6	Composition of Foods to be Fried and Fried Foods During Deep-Fat Frying	31
2.3.5.7	Filtration of Oils and Fats	32
2.3.6	Improvement of the Quality and Stability of Oils During Frying	32
2.3.7	Previous Studies on Canola and Sunflower Oils and their Blends with Palm Olein During Frying	34
2.4	Microwave Heating	36

2.4.1	Mechanism of Microwave Heating	38
2.4.2	Factors Affecting Dielectric Properties of Food	40
2.4.3	Effects of Frequency and Temperature	41
2.4.4	Effect of Moisture Content	42
2.4.5	Penetration Depth of Microwaves	43
2.4.6	Previous Studies on Canola and Sunflower Oils and their Blends with Palm Olein During Microwave Heating	44
3	METHODOLOGY	46
3.1	Food Materials and Chemicals	46
3.2	Experimental Methods	47
3.2.1	Frying Protocol	47
3.2.2	Microwave Heating	47
3.2.3	Analysis of the Oil Samples	48
3.2.3.1	Refractive Index	48
3.2.3.2	Free Fatty Acids	48
3.2.3.3	Iodine Value	49
3.2.3.4	Peroxide Value	50
3.2.3.5	<i>p</i> -Anisidine Value	50
3.2.3.6	Total Oxidation (TOTOX) Value	51
3.2.3.7	Oxidative Stability Index (OSI)	51
3.2.3.8	Viscosity	52
3.2.3.9	Specific Extinctions	53
3.2.3.10	Polymer Content	53
3.2.3.11	Total Polar Compound by Mini Column Method	54
3.2.3.12	Food Oil Sensor (FOS) Value	55
3.2.3.13	Color	55
3.2.3.14	Fatty Acid Composition by	

	GLC	55
	3.2.3.15 Thermal Analysis by DSC	56
	3.2.4 Statistical Analysis	57
4	RESULTS AND DISCUSSION (DEEP-FAT FRYING)	58
	4.1 Introduction	58
	4.2 Changes in Refractive Index (RI)	58
	4.3 Changes in Free Fatty Acid Content (FFA)	62
	4.4 Changes in Iodine Value (IV)	63
	4.5 Changes in Peroxide Value (PV)	64
	4.6 Changes in <i>p</i> -Anisidine Value (p-AV)	67
	4.7 Changes in Total Oxidation (TOTOX)	68
	4.8 Changes in Oxidative Stability Index (OSI)	68
	4.9 Changes in Viscosity	69
	4.10 Changes in Specific Extinctions at 233 and 269 nm ($E^{1\%}_{233}$ and $E^{1\%}_{269}$)	72
	4.11 Changes in Polymer Content (PC)	72
	4.12 Changes in Total Polar Compound (TPC)	73
	4.13 Changes in Food Oil Sensor (FOS) value	77
	4.14 Changes in Color	76
	4.15 Changes in Fatty Acids Composition (FAC)	79
	4.16 Changes in Thermal Properties	86
5	RESULTS AND DISCUSSION (MICROWAVE HEATING)	93
	5.1 Introduction	93
	5.2 Changes in Refractive Index (RI)	93
	5.3 Changes in Free Fatty Acid content (FFA)	96
	5.4 Changes in Iodine Value (IV)	96
	5.5 Changes in Peroxide Value (PV)	97
	5.6 Changes in <i>p</i> -Anisidine Value (p-AV)	100
	5.7 Changes in Total Oxidation (TOTOX)	100

5.8	Changes in Specific Extinctions at 233 and 269 nm ($E_{233}^{1\%}$ and $E_{269}^{1\%}$)	101
5.9	Changes in viscosity	104
5.10	Changes in Polymer Content (PC)	104
5.11	Changes in Total Polar Compound (TPC)	107
5.12	Changes in Food Oil Sensor (FOS) value	107
5.13	Changes in Fatty Acids Composition (FAC)	108
6	GENERAL CONCLUSIONS AND RECOMMENDATIONS FOR FUTURE WORKS	115
6.1	General Conclusions	115
6.2	Recommendations for Future Works	117
	REFERENCES	118
	Appendices A – E	133 – 149

LIST OF TABLES

TABLE NO.	TITLE	PAGE
2.1	World Palm Oil Production	10
2.2	Fatty acid composition of palm olein	11
2.3	World production rate of canola oil	14
2.4	Composition of Canola Oil	14
2.5	World production rate of sunflower oil	16
2.6	Fatty acids composition of sunflower oil	17
4.1	Refractive index, FFA and I_2 values of palm olein, canola oil, and their blends during frying	60
4.2	Refractive index, FFA and I_2 value of palm olein, sunflower oil, and their blends during frying	61
4.3	Peroxide, <i>p</i> -Anisidine, TOTOX and OSI values of palm olein, canola oil, and their blends during frying	65
4.4	Peroxide, <i>p</i> -Anisidine, TOTOX and OSI values of palm olein, sunflower oil, and their blends during frying	66
4.5	Viscosity, specific extinctions and polymer content of palm olein, canola oil, and their blends during frying	70
4.6	Viscosity, specific extinctions and polymer content of palm olein, sunflower oil, and their blends during frying	71
4.7	Total polar compound, FOS value and color of palm olein, canola oil, and their blends during frying	75
4.8	Total polar compound, FOS value and color of palm olein, sunflower oil, and their blends during frying	76
4.9	Fatty acids composition (%) of palm olein, canola oil, and their blends during frying	81
4.10	Fatty acids composition (%) of Sunflower Oil and its blends	

	during frying	82
4.11	Thermal Properties of Palm Olein, Canola, Sunflower and their blends during frying	88
5.1	Refractive index, FFA and I ₂ values of palm olein, canola oil, and their blends during microwave heating	94
5.2	Refractive index, FFA and I ₂ value of palm olein, sunflower oil, and their blends during microwave heating	95
5.3	Peroxide, <i>p</i> -Anisidine, TOTOX and OSI values of palm olein, canola oil, and their blends during microwave heating	98
5.4	Peroxide, <i>p</i> -Anisidine, TOTOX and OSI values of palm olein, sunflower oil, and their blends during microwave heating	99
5.5	Specific extinction and viscosity of palm olein, canola oil, and their blend during microwave heating	102
5.6	Specific extinction and viscosity of palm olein, sunflower oil, and their blend during microwave heating	103
5.7	Total polar compound, FOS value and color of palm olein, canola oil, and their blends during microwave heating	105
5.8	Total polar compound, FOS value and color of palm olein, sunflower oil, and their blends during microwave heating	106
5.9	Fatty acids composition (%) of palm olein, canola oil, and their blend during microwave heating	110
5.10	Fatty acids composition (%) of palm olein, sunflower oil, and their blend during microwave heating	111

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
2.1	Palm Seed	11
2.2	Canola Seeds	14
2.3	Sunflower Seed	17
2.4	Physical and chemical changes of oil during deep-fat Frying	22
2.5	The initiation, propagation, and termination of thermal oxidation of oil	26
2.6	Acyclic polymer formation from oleic acid during deep-fat frying	28
2.7	The interaction of microwaves with different materials	37
2.8	Contributions of various mechanisms to the loss factor of moisture materials as a function of frequency and temperature	40
2.9	Definition of penetration depth of microwave in a glossy material	43
4.1	Gas chromatogram of fresh PO used in frying experiment	83
4.2	Gas chromatogram of fresh CO used in frying experiment	83
4.3	Gas chromatogram of fresh SFO used in frying experiment	84
4.4	Gas chromatogram of fresh PC1 used in frying experiment	84
4.5	Gas chromatogram of fresh PC2 used in frying experiment	85
4.6	Gas chromatogram of fresh PSF1 used in frying experiment	85
4.7	Gas chromatogram of fresh PSF2 used in frying experiment	86
4.8	DSC cooling and heating thermograms of the fresh PO	89

4.9	DSC cooling and heating thermograms of the fresh CO	89
4.10	DSC cooling and heating thermograms of the fresh SFO	90
4.11	DSC cooling and heating thermograms of the fresh PC1	90
4.12	DSC cooling and heating thermograms of the fresh PC2	91
4.13	DSC cooling and heating thermograms of the fresh PSF1	91
4.14	DSC cooling and heating thermograms of the fresh PSF2	92
5.1	Gas chromatogram of fresh PO used in Microwave heating operation	112
5.2	Gas chromatogram of fresh CO used in Microwave heating operation	112
5.3	Gas chromatogram of fresh SFO used in Microwave heating operation	113
5.4	Gas chromatogram of fresh PC2 used in Microwave heating operation	113
5.5	Gas chromatogram of fresh PSF2 used in Microwave heating operation	114

LIST OF ABBREVIATIONS AND SYMBOLS

⁰ C	-	Degree Celsius
⁰ F	-	Degree Fahrenheit
ADC	-	Sunflower Oil
ANOVA	-	Analysis of Variance
cm	-	centimetre
CO	-	Canola Oil
cP	-	centipoise
DSC	-	Differential Scanning Calorimeter
g	-	grams
GLC	-	Gas Liquid Chromatography
h	-	Hour
HDL-C	-	High Density Lipoprotein- Cholesterol
kg	-	kilogramme
LDL-C	-	Low Density Lipoprotein- Cholesterol
Lp	-	Lipo- protein
min	-	minutes
ml	-	millilitre
mm	-	millimetre
MUFA	-	Monounsaturated Fatty Acid
nm	-	nanometer
PC1	-	Palm Olein: Canola Blend (20:80)
PC2	-	Palm Olein: Canola Blend (40:60)
PO	-	Palm Olien
PSF1	-	Palm Olein: Sunflower Oil (20:80)
PSF2	-	Palm Olein: Sunflower Oil (40:60)
PUFA	-	Polyunsaturated Fatty Acid
TFA	-	Total Fatty Acid

USFA	-	Unsaturated Fatty Acid
v/v	-	volume/ volume
µg	-	micrograms

LIST OF APPENDICES

APPENDIX	TITLE	PAGE
A	GLC Graphs for deep-fat frying (5th day)	134
B	DSC Graphs for Deep-Fat Frying (5th Day)	138
C	GLC Graphs for Microwave heating (20 minutes)	142
D	Apparatus	145

CHAPTER 1

INTRODUCTION

1.1 Background

Edible oils are used in cooking, baking and frying of foods. Consumption of edible oils has grown with the increase in the world population. The increase of health awareness and consciousness among consumers made the food industry more discriminating in the type of oil they use for food applications. Most native oils and fats have limited application in their unmodified forms, imposed by their tri-glyceride and fatty acids composition. By changing the natural physical and chemical characteristics of a fat or oil, it offers greater functionality for a large number of product formulations. Fats and oils can be modified through blending, fractionation, hydrogenation or combination of these processes (Petrauskaite *et al.*, 1998). Blending has long been used to modify oils and fats to improve their functionalities and thus optimizes their application in food products. It modifies the physicochemical properties of the oils, but without changing their chemical composition (Chu and Kung, 1997). Blending is also an economical way of producing health edible oils. Blending polyunsaturated oil with more saturated or monounsaturated oils, is an option to adjust fatty acid levels to optimal level (Nor Aini *et al.*, 1996).

Deep-fat frying is one of the oldest and most popular methods of preparation of food. Deep-fat frying is a process of immersing foods into hot edible oil or fat at temperature above boiling point of water. The oil is mostly heated to a temperature between 170 to 190°C which leads to heat and material transfer from the oil to the

food, resulting in cooking of the food; however, some of the oil is absorbed by the fried foods, and hence become a source of flavor, smell, taste and nutrition. Furthermore, undesirable off- flavor occur if deteriorated oil is used or if the oil is used over a longer period of time, as a result of the heat treatment a lot of deteriorative chemical processes take place, resulting in different changes of the oil. In this process, not only undesirable components are formed, but also compounds with adverse nutritional effects and potential hazards to humans health (Billek, 1985; Billek, 1992). Moreover, these changes may result in a decrease of quality and stability of the fried product. During frying, a number of reactions occur in the frying oil causing oxidative and hydrolytic degradation and polymerization of the oil. The main causes of oil degradation are oxidation, thermal treatment, and oil food interaction at high temperature (Gary *et al.*, 1997). The quality of the oil used in deep-fat frying contributes to the quality of the fried food. The oil or fat used in the frying operation becomes part of the food we eat and is, of course, the major factor in the quality and nutritional value of the food we eat. In addition, highly oxidized oils may also produce polycyclic aromatic hydrocarbons that are known as carcinogens (Sebedio *et al.*, 1990; Romero *et al.*, 1998). On the other hand, microwave heating is a widespread procedure for food preparation and manufacturing, commonly employed in both domestic and catering operations for its high speed and convenience, as compared to conventional heating treatments (Burfoot *et al.*, 1990). Heating of food by microwave outcomes from the molecular friction of electric dipoles, under an oscillating electric field of specific frequency (Mudgett, 1989). Advantages of microwave heating include savings in time and energy and easy to use. These advantages make it one of the most attractive cooking methods.

Palm olein is a highly monounsaturated oil, which is rich in oleic and is currently said to be oxidative stable acids (Nor Aini *et al.*, 1993). Besides being marked as liquid oil, palm olein can be promoted for blending with other edible oils (Lin, 2002). It is being used increasingly in frying operations, and because of its inherent excellent frying properties, improve the frying quality of other vegetable oils when it is blended with them (Kun, 1990). Canola oil is the best source of omega 3 fatty acids which helps to reduce cholesterol in the blood stream. It contains the lowest concentration of saturated fatty acids (7%) among vegetable oils (Ackman, 1990). Canola oil, because of its high content of PUFA it is considered superior to

many vegetable oils, but inferior in terms of thermal stability at high temperatures (Malcolmson *et al.*, 1994). Sunflower oil is one of the healthiest and popular oil in the world. It is also excellent household oil, great for both frying and baking. It is a rich source of vitamin E and has a high level of linoleic acid (Huang *et al.*, 1981; Beddows, 2000).

In view of the foregoing it has become paramount to undertake the necessary measurements by a combination of the most effective analytical and instrumental analysis to evaluate the quality and stability of canola and sunflower oils and their blends with palm olein during frying and microwave heating.

1.2 Problem Statement

Most healthy vegetable oils like canola oil, sesame oil, sunflower oil, corn oil etc. are expensive and cannot be reuse for so many number of frying due to the formation of oxidation and polymerization products after subjecting to high temperature and air, which makes it difficult for the food processing companies, restaurant and the consumers to purchase. In fact, the applications of canola and sunflower oils in house hold and in food industry are limited due to their high cost. Today, palm oil and its products, mainly palm olein, belong to the most important oils used for the preparation of fried food. The reason is that the oil is relatively cheap, it is available in huge amounts, it has a high oxidative stability and results in high-quality and tasty foods. However, the price is an important modulatory factor on the market, because for the production of fried food the oil is the most expensive part. Addition of oxidation-stable palm olein to canola and sunflower oils may affect the oxidative stability of canola and sunflower oils during frying or microwave heating. So, blending is the perfect way of obtaining qualitative and stable oils. Blending canola and sunflower oil with palm olein may lead to nutritious, qualitative and stable oils. Furthermore, blending canola and sunflower oil with palm olein will reduce the cost of purchase of the vegetable oils.

Secondly, the food industry and public health authorities are worried and concerned in relation to potential health hazards generated by the consumption of

oxidized products, which most often contain lipid polymers resulting from oils used repeatedly in frying processes. The stability of the oil is also very important, because stability to oxidation during a prolonged exposure to high temperature is one of the main characteristics that industrial and domestic frying oils should possess, as its physical, chemical and nutritional properties may undergo significant modification during thermo-oxidation. From a practical point of view, commercial and industrial frying oil operators have no idea on when to discard frying oils. They want to know when the frying oils should be discarded, in other words when the frying oil needs to be dumped. Unfortunately, there is no simple answer to such question. A specific method may be ideal for one operation but useless in another. The end point of frying oil is therefore, dependent on mainly analytical measurements, good judgment and knowledge of the particular frying operation and the type of frying oils. This research work may help in consumer awareness on when to discard the oils after several uses and also help in buying the healthy oil from the market. On the other hand, amongst the various cooking techniques, microwave oven heating is a more recent development than other traditional cooking techniques. Due to its advantages (time and energy savings), this technique has become one of the most attractive cooking methods. This justifies its increasing use for studying the heating behavior of various foodstuffs.

Therefore, proper blending of canola and sunflower oils with palm olein in the optimal proportion may results in production of oil blends with improved stability, nutritional value and flavor characteristics.

1.3 Objectives of study

- To evaluate the changes in quality and thermooxidative degradation of highly unsaturated canola and sunflower oils and their blends with palm olein during frying of potato pieces and microwave heating.
- To produce blended oils that are less costly with good stability which can withstand domestic cooking and commercial frying or microwave heating operations.

1.4 Scopes of Study


To achieve the objectives of this research work, the following scopes were identified:

- To evaluate the impacts of addition of palm olein at different concentrations on the frying performance of canola oil and sunflower oil.
- To determine how microwave heating may affect the quality and stability of canola and sunflower oils and their blends with palm olein.
- To attempt to link the oil's deterioration to fatty acid composition during frying or microwave heating.

1.5 Significance of Study

The quality and stability of oils is very important to the human diet, to avoid consumption of unnecessary fats which can affect their body system and health. Moreover, the advantage of using blend as a means of modifying oils is that it is easier to use and less costly. Blending also serves to improve and enhance the nutritional and functional qualities of the oils, by combining the good attributes of the two blended oils into one. This research work may serve as a milestone towards development of effective frying operations and may increase consumer awareness toward nutritional values and health effects. This research can results in oil blends which could meet nutritional needs with improved stability for domestic and industrial deep fat frying. This study will also help to understand on how heat treatments by microwave oven may affect the oxidative degradation of canola and sunflower oils and their blends with palm olein.

1.6 Work Flow Chart



1.7 Thesis Outlines

The thesis is organized as follows:

In chapter 2, a brief description on palm olein, canola and sunflower oils was mentioned. The theory and application of heat transfer during deep-fat frying and the mechanism of microwave heating are also presented in this chapter. Moreover, previous studies on deep-fat frying and microwave heating of canola and sunflower oils and their blends with palm olein are reviewed. The effect of oil quality during frying and the method on how to improve the quality of the oils are highlighted.

In chapter 3, the experimental setups and procedures for frying and microwave heating employed in this study are described. The methodologies of each and every experiment considered in this study, are described in detailed.

Chapter 4 presents the results and discussion of deep-fat frying of the samples obtained from both instrumental and analytical methods.

Chapter 5 presents the results and discussion of microwave heating of the samples obtained from both instrumental and analytical methods.

Finally, chapter 6 is the concluding chapter. This chapter summarizes and concludes the research that has been carried out in this study. Future works are also suggested at the end of this chapter.

REFERENCES

- Abdulkarim, S. M., Long, K., Lai, O. M., Muhammad, S. K. S., Ghazali, H. M., (2007). Frying quality and stability of high-oleic *Moringa oleifera* seed oil in comparison with other vegetable oils. *Food Chem.* **105**: 1382–1389.
- Abdulkarim, S. M., Myat, M. W., Ghazali, H. M., Roselina, K. and Abbas, K. A. (2010). Sensory and physicochemical qualities of palm olein and sesame seed oil blends during frying of banana chips. *J. Agric. Sci.* **2**: 18-29.
- Ackman, R.G. (1990). *Canola Fatty Acids- an ideal Mixture for Health, Nutrition and food use*. In: Canola and repeseed. Eds. Shahidi, F. and Reinhold, V.N., New York, pp. 81-89.
- AHA. (2011). American heart association. Retrieved on the 27th December 2011 from <http://www.heart.org/HEARTORG/>
- Albi, T., Lanzon, A., Guinda, A., Pérez-Camino, M. C. and Leon, M. (1997). Microwave and conventional heating effects on some physical and chemical parameters of edible fats. *J. Agric. Food Chem.* **45**: 3000-3003.
- Alireza, S., Chin, P. T., Hamad, and Yaakob, C. M. (2010). Effect of Vegetable based Oil blends on physicochemical properties of Oils during Deep Fat Frying. *Am. J. Food Tech.* **5**: 310-323.
- Al-Kahtani, H., (1991). Survey of quality of used frying oils from restaurants. *J. Am. Oil Chem. Soc.* **68**: 857–862.
- Allman-Farinelli, M.A., Gomes K., Favaloro, E.J, Petocz, P. (2005). A diet rich in high-oleic acid sunflower oil favorably alters low-density lipoprotein cholesterol, triglycerides, and factor VII coagulant activity. *J. Am. Diet Assoc.* **105**: 1071-1079.
- AOCS. (1987). *Official Methods and Recommended Practices of the American Oil Chemists' Society*. 4th ed., AOCS press, Champaign.

- APOC. (2004). American palm oil council. Retrieved on July 7th 2011 from <http://www.americanpalmoil.com>
- APOC. (2008). American palm oil council. Retrieved on July 7th 20011 from <http://www.americanpalmoil.com>
- Ashraf, G. A., Reza, F. and Mohammad, H.H.K. (2010). Frying stability of canola oil in presence of pumpkin seed and olive oils. *Eur. J. Lipid Sci. Technol.* **112**: 871-877.
- Ayappa, K.G., Davis, H.T., Crapiste, G., Davis, E.A. and Gordon, J. (1991). Microwave heating, an evaluation of power formulations. *J. Chem. Eng. Sci.* **46**: 1005-1016.
- Beddows, C. B. (2000). Preservation of alpha-tocopherol in sunflower oil by herbs and spices. *Int. J. Food Sci. Nutr.* **51**: 327-340.
- Bertrand, M. (2007). Use of palm oil for frying in comparison with other high-stability oils. *Eur. J. Lipid Sci. Technol.* **109**: 400-409.
- Billek, G. (1985). *Heated Fats in the Diet*. In: The Role of Fats in Human Nutrition. Eds. Padley, F. B., Podmore, U., Horwood, E., Horwood, E., West Sussex, U.K., pp. 163-171.
- Billek, G. (1992). Die veränderungen von nahrungsfettenbeihöheren temperature. *Fat. Sci. Technol.* **94**: 161-172.
- Billek, G., Guhr, G., Waibel, J., (1978). Quality assessment of used frying fats: A comparison of four methods. *J. Am. Oil Chem. Soc.* **55**: 728-732.
- Blumenthal, M. M., (1996). *Frying Technology*. In: Edible Oil and Fat Products: Products and Application Technology. In: Hui, Y. (Ed.), *Bailey's Industrial Oil & Fat Products*, 5th edn., Vol. 3., John Wiley and Sons, Inc., N.Y, pp. 429–481.
- Blumenthal, M.M. (1991). A new look at the chemistry and physics of deep-fat frying. *J. Food Technol.* **45**: 68-71.

- Boskou, G., Salta, F.N., Chiou, A., Troullidou, E. and Andrikopoulos, N.K. (2006). Content of Trans, trans-2, 4-decadienal in Deep-Fried and Pan-Fried Potatoes. *Eur J. Lipid Sci. Technol.* **108**: 109-115.
- Buffler, C. and M. Stanford. (1991). The effects of dielectric and thermal properties on the microwave heating of foods. *Microwave World* **9**: 15-24.
- Burfoot, D., James, S.J., Foster, A.M., Self, K.P., Wilkins, T.J. and Philips, I. (1990). In: Temperature Uniformity After Reheating in Microwave Ovens. Eds. Field, R.W. and Howell, J.A. *Processing engineering in the food industry*, Elsevier Applied Science, New York, **2**: 1-14.
- Burfoot, D., James, S.J., Foster, A.M., Self, K.P., Wilkins, T.J. and Philips, I. (1990). *Temperature Uniformity After Reheating in Microwave Ovens*. In: R.W. Field, and J.A. Howell Eds., *Processing engineering in the food industry*, Vol. 2, Elsevier Applied Science, New York, pp. 1-14.
- CCC. (1990). Canola Council of Canada. Canola oil and meal: standards and regulations. Winnipeg, Manitoba: Canola Council of Canada Publication. p. 4.
- CCC. (2008). Canola Council of Canada. Comparison of Dietary Fats Chart. Retrieved on July 25th 2011 from www.canola-council.org.
- CCC. (2009). Canola Council of Canada. Canadian Canola Industry: Overview of Canada's Canola Industry. Retrieved on June 28th, 2011 from http://www.canolacouncil.org/ind_overview.aspx.
- Che Man, Y. B., Liu, J. L., Jamilah, B., Abdul Rahman, R. (1999). Quality changes of refined-bleached-deodorized (RBD) palm olein, soybean oil and their blends during deep-fat frying. *J. Food Lipid.* **6**: 181-193.
- Che Man, Y. B., Hussin, W. R., (1998). Comparison of the frying performance of refined, bleached and deodorized palm olein and coconut oil. *J. Food Lipids.* **5**:197–210.
- Che Man, Y. B., Liu, J. L., Jamilah, B., Rahman, R. A., (1999). Quality changes of refined-bleached-deodorized (RBD) palm olein, soybean oil and their blends during deep-fat frying. *J. Food Lipids* **6**:181–193.

- Che Man, Y. B., Swe, P. Z., (1995). Thermal analysis of failed-batch palm oil by differential scanning calorimetry. *J. Am. Oil Chem. Soc.* **72**: 1529–1532.
- Choe, E. and Min, D. B. (2007). Chemistry of lipids and deep fat frying oils. *J. Food Sci.* **72**:77-86
- Chu, Y. H., Kung, Y. H. (1997). A study on oxidative stability of vegetable oil blends. *Ship in Kexue* **24**: 389-397.
- Chung, J., Lee, J. and Choe, E. (2004). Oxidative stability of soybean and sesame oil mixture during frying of flour dough. *J. Food Sci.* **69**: 574-578.
- Clark, D. E., Folz, D.C., and West, J. K. (2000). Processing material with microwave energy. *J. Mat. Sci. Eng.* **287**: 153-158.
- Clark, W.L. and Serbia, G.W. (1991). Safety aspects of frying fats and oils. *J. Food Technol.* **45**: 84-89.
- Costa, R.M., Oliveira, F.A.R., Delaney, O., and Gekas, V. (1999). Analysis of the heat transfer coefficient during potato frying. *J. Food Eng.* **39**: 293-299.
- Cuesta, C., Sanchez-Muniz, F. J., Garrido-Polonio, C., Lopez-Varela, S. and Arroyo, R. (1993). Thermo oxidative and hydrolytic changes in sunflower oil used in frying with a fast turnover of fresh oil. *J. Am. Oil Chem. Soc.* **70**:1069-1073.
- Cuesta, C., Sanchez-Muniz, F. J., Hernandez, I., (1991). Evaluation of nonpolar methyl esters by column and gas chromatography for the assessment of used frying olive oils. *J. Am. Oil Chem. Soc.* **68**: 443–445.
- Darnoko, D., Cheryan, M., (2000). Kinetics of palm oil transesterification in a batch reactor. *J. Am. Oil Chem. Soc.* **77**:1263–1267.
- Datta, A. K. and Hu, W. (1992). Optimization of quality in microwave heating. *J. Food Technol.* **46**: 53-56.
- De Marco, E., Savarese, M., Parisini, C., Battimo, I., Falco, S. and Sacchi, R., (2007). Frying performance of a sunflower/ palm oil blend in comparison with pure palm oil. *Eur. J. Lipid Sci. Technol.* **109**: 237–246.

- Decareau, R. V. (1985). *Microwaves in the Food Processing Industry*. Academic Press, Inc., New York.
- Dobarganes, C., Marquez-Ruiz, G. and Velasco, J. (2000). Interactions between fat and food during deep-frying. *Eur J, Lipid Sci Technol*. **102**: 521-528.
- Dostálová, J., Hanzlík, P., Réblová, Z., and Pokorný, J. (2005). Oxidative changes of vegetable oils during microwave heating. *Czech J. Food Sci*. **23**: 230-239.
- Emma Chiavaro, Maria Teresa Rodriguez-Estrada, Elena Vittadini and Nicoletta Pellegrini. (2010). Microwave heating of different vegetable oils: Relation between chemical and thermal parameters. *J. Food Sci. Technol*, **43**: 1104-1112.
- Enríquez-Fernández, B. E., Álvarez de la, C. Y. Y. L., and Sosa-Morales, M. E., (2011). Comparison of the stability of palm olein and a palm olein/canola oil blend during deep-fat frying of chicken nuggets and French fries. *Int. J. Food Sci. Technol*. **46**: 1231–1237.
- Farag, R. S., Hewedi, F. M., Abu-Raiia, S. H. and Elbaroty, G. S. (1992). Comparative study on the deterioration of oils by microwave and conventional heating. *J. Food Protect*. **55**: 722-727.
- Farhoosh, R., Kenari, R. E., Poorazrang, H., 2009. Frying stability of canola oil blended with palm olein, olive, and corn oils. *J. Am Oil Chem Soc*. **86**: 71–76.
- Farkas, B. E.(1994). *Modeling immersion frying as a moving boundary problem*. Ph.D. Dissertation, University of California, Davis.
- Fatihanim M. N, Mohamed S, Nor Aini I, Razali I. (2008). Antioxidative properties of Pandanus amaryllifolius leaf extracts in accelerated oxidation and deep frying studies. *J. Food Chem*. **110**: 319-327
- Fereidoon S. (2005). Bailey's Industrial Oil and Fats Products: Edible Oil and Fat Product, Chemistry, Properties and Health Effects. USA: Wiley inter science, pp. 244-245
- Fick G. N, (1989). *Sunflower*. In: *Oil Crops of the World*. Ed. Robbelen, G., Downey, R. K. and Ashri, A. McGraw-Hill, New York, pp. 301-318.

- Frankel, E. N., and S.W. Huang, (1994). Improving the Oxidative Stability of Polyunsaturated Vegetables Oils by Blending with High-Oleic Sunflower Oil. *J. Am Oil Chem. Soc.* **71**: 255-259.
- Frankel, E. N., K. Warner, and K. J, (1985). Moulton, Effects of Hydrogenation and Additives on Cooking Oil Performance of Soybean Oil. *J. Am Oil Chem. Soc.* **62**: 1354-1358.
- Frankel, E. N., Smith, L. M., Hamblin, C. L., Creveling, R. K. and Clifford, A. J. (1984). Occurrence of cyclic fatty acid monomers in frying oils used for fast foods. *J. Am. Oil Chem. Soc.* **61**: 87-90.
- Frega, N., Mozzon, M. and Lecker, G. (1999). Effects of free fatty acids on oxidative stability of vegetable oil. *J. Am. Oil Chem. Soc.* **76**: 325-329.
- Fristch, C. W. Ž. (1981). Measurements of frying fat deterioration: a brief review. *J. Am. Oil Chem. Soc.* **58**: 272-274.
- Gary, R. T., Gerhard, H. F., Lan, T. D., (1997). Effect of heating on the characteristics and chemical composition of selected frying oils and fats. *J. Agri. Food Chem.* **45**: 3244–3249.
- Giese, J. H. (1992). Special report, in advances in microwave food processing. *J. Food Technol.* **46**: 118-123.
- Goldblith, S.A. (1967). Basic principles of microwaves and recent developments. *Adv. Food Res.* **15**: 277–301.
- Gopal, J. (2001) *The Development of Malaysia's Palm Oil Refining Industry: Obstacles, Policy and Performance*. Ph.D. Thesis, Imperial College, London.
- Gordon, M. H and Kourimska, L. (1995). The effects of antioxidants on changes in oils during heating and deep frying. *J. Sci. Food Agric.* **68**: 347-353.
- Gray, J.I., Measurements of Lipid Oxidation. A Review, *J. Am. Oil Chem. Soc.* **55**:539–546 (1978).
- Graziano, V. J., (1979). Portable instrument rapidly measures quality of frying fat in food service operations. *Food Technol.* **33**:50–57.

- Harcharan Singh Khera (1976). Oil Palm Industry: An Economic Study, Kuala Lumpur: University of Malaya Press.
- Harish K. S, Barinder K, Bhavesh C. S and Charanjiv S. (2006). Thermal behavior of pure rice bran oil, sunflower oil and their model blends during deep fat frying. *J. fats and oils*, **57**: 376- 381.
- Hartley, C. W. S. (1988). The Oil Palm, 3rd edn. Longman Scientific and Technical, Harlow, U.K.
- Heiser C. B, (1976). Sunflowers: *Helianthus (Compositae– Heliantheae)*, in *Evolution of Crop Plants*, ed. by Simmonds NW. Longmans Green, London, pp. 36–38.
- Houhoula, D. P., Oreopoulou, V. and Tzia, C. (2003). The Effect of Process Time and Temperature on the Accumulation of Polar Compounds in Cottonseed Oil during Deep-fat Frying. *J. Sci. Food Agric.* **83**: 314-319.
- Huang, A.S., Hsieh, O.A.L., Haung, C.L. and Chang, S.S. (1981). A comparison of the stability of sunflower oil and corn oil. *J. Am. Oil Chem. Soc.* **58**: 997-1001.
- Innawong, B. (2001). *Improving fried product and frying oil quality using nitrogen gas in a pressure frying system*. Ph. D. Dissertation, Virginia Polytechnic Institute and State University.
- Khan, M. I., Asha, M. R., Bhat, K. K., Khatoon, S., (2011). Studies on chemical and sensory parameters of coconut oil and its olein blends with sesame oil and palm olein during wheat flour-based product frying. *J. Food Sci. Technol.* **48**:175–182.
- Kim, I. H., Kim, C. J., Kim D. H. (1999). Physicochemical properties of methyl linoleate oxidized at various temperatures. *Korean J. Food Sci. Technol.* **31**: 600-605.
- Kimber, D. S, and Mc Gregor D. I. (1995). *The Species and Their Origin, Cultivation and World Roduction*. In: Ed. Kimber D.S. and McGregor D. I., Brassica oilseeds: production and utilization. Wallingford, England: CAB Intl. p 1-7.

- Kochhar, S. P. and Gertz, C. (2004). New theoretical and practical aspects of the frying process. *Eur. J. Lipid Sci. Technol.* **106**: 722-727.
- Kochhar, S. P., (2001). *The composition of frying oils*. In: Rossell J. B. (Ed.), *Frying – Improving Quality*. Wood head Publishing Ltd., Cambridge UK.
- Kress Rogers, E., Gillat, P. N., Rosel, J. B., (1990). Development and evaluation of a novel sensor for the *in situ* assessment of frying oil quality. *Food Control*. **13**:163–178.
- Kuang, W. and S. O. Nelson. (1998). Low-frequency dielectric properties of biological tissues: a review with some new insights. *Trans. ASAE* **41**:1 73-184.
- Kun, T.Y. (1990). Improvements in the frying quality of vegetable oil by blending with palm olein. palm oil development, No 15, PORIM, Bangi, Selangor, Malaysia
- Lascaray, L. (1949). Mechanism of fat splitting. *Ind. Eng. Chem.* **41**:786-90.
- Lawson, H. (1995). *Food Oils and Fats: Technology, Utilization and Nutrition*. New York: Chapman and Hall, pp. 15-20.
- Leyla, T., Mehmet, S. A., Emin, Y., (2009). Physicochemical changes in hazelnut, olive pomace, grapeseed and sunflower oils heated at frying temperatures. *J. Food Sci. Technol. Res.* **15**:519–524.
- Lie-Ken-Jie, M. and Yan-Kit, C. (1988). The use of a microwave oven in the chemical transformation of long chain fatty acid esters. *J. Lipids*. **23**: 367-369.
- Lin, S.W. (2002). *Vegetable Oils in Food Technology, Composition, Properties and Uses*. In: Palm oil. Eds. Gunstone, F.D., SRC Press LLC, pp. 59-86.
- Liu, H., and P. J, (1992). White, High-Temperature Stability of Soybean Oils with Altered Fatty Acid Compositions, *J. Am. Oil Chem. Soc.* **69**: 533-537.
- Lukešová, D., Dostálová, J. E., El-Moneim, M. and Svárovská, M. (2009). Oxidation Changes of Vegetable Oils during Microwave Heating. *J. Food Sci.* **27**: S178-S181.

- Lusas, E.W. (1985). *Sunflower Seed Protein*. In: New Protein Foods. Eds. Altschul, A.M. and Wilcke, H, L. Academic Press, pp. 393–433.
- Malaysia (1975) Third Malaysia Plan 1976-1980, Kuala Lumpur: Government Printers. 5 MPOB 2000. "History of Oil Palm in Malaysia", Unpublished mimeo, Malaysian Oil Palm Board, Kuala Lumpur.
- Malcolmson, L.J., Vaisey-Genser, M., Przybylski, R. and Eskin, N.A.M. (1994). Sensory stability of canola oil: present status of shelf life studies. *J. Am. Oil Chem. Soc.* **71**: 435-440.
- Mariod, A., Matthäus, B., Eichner, K., Hussein, I. H., (2006). Frying quality and oxidative stability of two unconventional oils. *J. Am. Oil Chem. Soc.* **83**:529–538.
- Maskan, M. (2003). Change in color and rheological behavior of sunflower seed oil during frying after adsorbent treatment of used oil. *Eur. J. Food Res. Technol.* **218**: 20-25.
- Matthaus, B., (2006). Utilization of high-oleic rapeseed oil for deep fat frying of French fries compared to other commonly used edible oils. *Eur. J. Lipid Sci. Technol.* **108**: 200–211.
- Mazza, G. and Qi, H. (1992). Effect of after-cooking darkening inhibitors on stability of frying oil and quality of French fries. *J. Am. Oil Chem. Soc.* **69**: 847-853.
- MCGA. (2008). Manitoba Canola Growers Association. Retrieved on July 25th 2011 from www.mcgacanola.org.
- Metaxas, A. C. and R.J. Meredith (1993). *Industrial Microwave Heating*. Peter Peregrinus Ltd., London.
- Min, D.B. and Boff, J.M. (2002). *Lipid Oxidation of Edible Oil*. 2nd ed., Marcel Dekker, New York.
- Minar, M. H., Safinaz, M. E. and Hassan E, M. (2003). Changes occurring in vegetable oils composition due to microwave heating. *Grasas y Aceites*, **54**: 343-349.

- Moreira, R.G., Sun, X., Chen, Y. (1997). Factors affecting oil uptake in tortilla chips in deep-fat frying. *J Food Eng.* **31**:485-98.
- Moulton, K. J., Beal, R. E. Warner, K. and Boundy, B. K. (1975). Flavor Evaluation of Copper–Nickel Hydrogenated Soybean Oil and Blends with Unhydrogenated Oil. *J. Am Oil Chem. Soc.* **52**: 469-472.
- MPOB. (2010). Malaysian Palm Oil Council. Retrieved on the 16th September 2011 from <http://www.mpoc.org.my/>
- Mudgett, R.E. (1989). Microwave food processing. *J. Food Technol.* **43**: 117-126.
- Nawar W. W. (1984). Chemical changes in lipids produced by thermal processing. *J. Am. Oil Chem. Soc.* **61**: 299-302.
- Naz, S., Siddiqi, R., Sheikh, H. and Sayeed, S.A. (2005). Deterioration of Olive, Corn, and Soybean Oils due to Air, Light, Heat, and Deep-Frying. *Food Res. Intl.* **38**: 127-134.
- Nelson, S. O., &Kraszewski, A. W. (1990). Grain moisture content determination by microwave measurements. *Trans. ASAE* **33**: 1303-1307.
- Noakes, M., Nestel, P.J. and Clifton, P.M. (1996).Commercial frying fats and plasma lipid-lowering potential. *Aus J. Nutri. Dietic.* **53**: 25-30.
- Nor Aini, I., Yusof, B., Hanirah,H. Rosilah, B. (1993). Effect of some additives on resistance to crystallization of palm oil. *Elaeis.* **5**: 47-64.
- Normand, L., Eskin, N. A. M., Przybylski, R., (2001). Effects of tocopherols on the frying stability of regular and modified canola oils. *J. Am. Oil Chem. Soc.* **78**: 369–373.
- NSA. (2012). National Sunflower Association. Retrieved on March 7th from <http://www.livestrong.com/article/82773-health-benefits-sunflower-il/#ixzz23L6C1Xpq>
- Oil world. (2011). Retrieved on 18th December 2011 from [http://www.poram.org.my/v1/poram/statistic/3.%20World%20PalmOil%20Pro](http://www.poram.org.my/v1/poram/statistic/3.%20World%20PalmOil%20Production.pdf)duction.pdf

- Onal, B., Ergin, G., (2002). Antioxidative effects of α -tocopherol and ascorbyl palmitate on thermal oxidation of canola oil. *Nahrung*, **46**:420–426.
- Oztop M. H, Sahin S., Sumnu, G. (2007). Optimization of microwave frying of potato slices by using Taguchi technique. *J. Food Eng.* **79**: 83-91
- Paul, S. and Mittal, G.S. (1997). Regulating the use of degraded oil/fat in deep-fat/oil food frying. *Crit. Rev. Food Sci. Nutr.* **37**: 635-62.
- Peled M., Gutfinger T., Letan A. (1975). Effect of water and BHT on stability of cottonseed oil during frying. *J. Sci. Food Agric.* **26**, 1655-1666.
- Perkins, E.G. and Erickson, M.D. (1996). *Regulation of Frying Fat and Oil in Deep Frying*. In: Chemistry, Nutrition, and Practical Application. Champaign I. L., ACCS Press, pp. 323 -334
- Petrauskaite, V., De Greyt, W., Kellens, M. and Huyghebaert, A. (1998). Physical and chemical properties of trans- free fats produced by chemical interesterification of vegetable oil blends. *J. Am. Oil. Chem. Soc.* **75**: 489-493.
- Pokorny, J. (1989). *Flavor Chemistry of Deep Fat Frying in Oil*. In: Eds. Min, D. B. and Smouse. T. H., Flavor chemistry of lipid foods. AOCS Press, Champaign Ill, pp. 113-115.
- PORIM (1995). *PORIM test methods*. Malaysia: Palm Oil Research Institute of Malaysia, Ministry of Primary Industries.
- Potter, N.N. (1986). Food science. Avi, Westport, Conn. USA, pp. 322-353.
- Premavalla, K. S., Madhura, C. V. and Arya, S.S. (1998). Storage and thermal stability of refined cottonseed oil: mustered blend. *J. Food Sci. Technol.* **35**: 530–532.
- Prevot, A., Desbordes, S., Morin, O. and Mordret, F. (1988). *Volatiles and Sensory Effects from Frying Oils*. In: Eds. Varela, G., Bender, A. E. and Morton, I.D. Frying of food: principles, changes, New Approaches. Ellis Harwood Ltd., UK, pp. 155-165.

- Purdy, R. H., and B. J. (1967). Campbell, High-Oleic Safflower Oil. *J. Food Tech.* **21**:349-352.
- Radwan, S.F, Magdy A. S. El-Agaimy, and Bothaina S. A. (2010). Effects of Mixing Canola and Palm Oils with Sunflower Oil on the Formation of Trans Fatty Acids during Frying. *J. Food Nutrition Sci.* **1**: 24-29.
- Razali, I. (2005). Palm oil and palm olein frying applications. *Asia Pac. J. Clin. Nutr.* **14**: 414-419.
- Reeves J. B, Weihrauch, J. L. (1979). Consumer and Food Economics Institute, *Composition of foods: fats and oils*. Agriculture handbook 8-4. Washington, D.C.: U.S. Dept. of Agriculture, Science and Education Administration. p. 4.
- Reza, F. E. K. and Hashem, P. (2008). Frying stability of canola oil blended with palm olein, olive and corn oils. *J. Am. Oil Chem. Soc.* **86**: 71-76
- Romero, A., Cuesta, C. and Sanchez-Muniz, F. J. (1998). Effect of oil replenishment during deep-fat frying of frozen foods in sunflower oil and high-oleic acid sunflower oil. *J. Am. Oil Chem. Soc.* **75**:161-167.
- Rossel, J.B., 1989. *Measurement of rancidity*. In J.C. Allen and R. J. Hamilton (Eds.), *Rancidity in foods*, England: Elsevier Science. pp. 45–49.
- Sanchez-Muniz, F. J., Cuesta, C., Lopez-Varela, M.C., Garrido-Polonio, M.C. and Arroyo, R. (1993). *Evaluation of the Thermal Oxidation rate of Sunflower Oil using various Frying Methods*. In: Ed. Apple W. T. H. *Proceedings of World Conference on Oil Seed and Technology and Utilization*. Champaign, Ill., AOCS Press, pp. 448-452.
- Sebedio, J. L. Bonpunt, A. Grandgirard, A. Prevost, J. (1990). Deep fat frying of frozen pre-fried French fries: influence of the amount of linolenic acid in the frying medium. *J. Agric. Food Chem.* **38**: 1862–1867.
- Seberry, D. E, Mailer R. J. and Parker P. A. (2008). The quality of Australian canola. Australia: Australian Oilseeds Federation, NSW Dept. of Primary Industries.

- Shahidi, F., Wanasundara, U. N., (2002). Methods for measuring oxidative rancidity in fats and oils. In: Akoh C. C., Min D. B. (Eds.), *Food lipids: Chemistry, nutrition, and biotechnology*, 2nd edn., Marcel Dekker, Inc. N. Y. pp. 465–482.
- Shahidi, F.; Wanasundara, U. N., (2002). *Methods for Measuring Oxidative Rancidity in Fats and Oils*. In: *Food lipids: Chemistry, nutrition, and biotechnology*, Eds. Akoh C.C. and Min D.B., 2nd ed., Marcel Dekker Inc., N. Y, pp 465-482.
- Sibel, K., Sebnem, S., (2011). Changes in total polar compounds, peroxide value, total phenols and antioxidant activity of various oils used in deep fat frying. *J. Am. Oil Chem. Soc.* **88**: 1361–1366.
- Siew, W. L., Ng, W. L., (1995). Diglyceride content and composition as indicators of palm oil quality. *J. Sci. Food Agric.* **69**:73–79.
- Stevenson, S. G., Vaisey-Genser, M. and Eskin, N. A. M. (1984). Quality Control in the use of Deep Frying Oils. *J. Am. Oil. Chem. Soc.* **61**: 1102-1108.
- Sumnu, G. (2001): A review on microwave baking of foods. *Int. J. Food Sci. Technol.* **36**: 117-127.
- Tabee, E., Jagerstad, M., Paresh, C. D., (2009). Frying quality characteristics of french fries prepared in refined olive oil and palm olein. *J. Am. Oil Chem. Soc.* **86**:885–893.
- Tan, C. P., Che Man Y. B. (1999). Differential scanning calorimetric analysis for monitoring the oxidation of heated oils. *J. Food Chem.* **67**: 177–184.
- Tan, C. P., Che Man, Y. B., Jinap, S. and Yusoff, M. S. A. (2002). Effects of microwave heating on the quality characteristics and thermal properties of RBD palm olein, *Innov. Food Sci. Emerg. Technol.* **3**: 157-163.
- Thais, M. F. S., Vieira, and Marisa, A. B., Regitanod, A. (2001). Canola oil thermal oxidation during oven test and microwave heating. *Lebensm.-Wiss. u.-Technol.* **34**: 215-221.

- Tompkins, C. and Perkins, E.G. (2000). Frying performance of low-linolenic acid soybean oil. *J. Am. Oil Chem. Soc.* **77**: 223-229.
- Tsaknis, J., Lalas, S. (2002). Stability during frying of *Moringa oleifera* seed oil variety “Periyakulam 1”. *J. Food Comp. Anal.* **15**:79–101.
- Tsuyuki, H. (1982). Utilization of high frequency and microwave heating in food industry. *J. Jpn. Soc. Food Sci. Technol.* **29**: 123-130.
- Tsuzuki, W., Matsuoka, A., Ushida, K. (2010). Formation of trans fatty acids in edible oils during the frying and heating process. *Food Chem.* **123**: 976–982.
- Tyagi, V. K. and Vasishtha, A. K. (1996). Changes in the characteristics and composition of oils during deep-fat frying. *J. Am. Oil Chem. Soc.* **73**: 499-506.
- Warner, K. & Mounts, T. L., (1993). Frying stability of soybean and canola oils with modified fatty acid compositions. *J. Am. Oil Chem. Soc.* **70**: 983–988.
- Warner, K., and Knowlton, S. (1997). Frying quality and oxidative stability of high-oleic corn oils. *J. Am. Oil Chem. Soc.* **74**: 1317-1321.
- Warner, K., Orr, P., Glynn, M., (1997). Effect of fatty acid composition of oils on flavour and stability of fried foods. *J. Am. Oil Chem. Soc.* **74**:347–356.
- Williams, G. M., Iatropoulos, M. J. and Whysner, J. (1999) Safety assessment to butylatedhydroxyanisole and butylatedhydroxytoluene as antioxidant food additives. *Food Chem Toxicol.* **37**:1027-1038
- Wilson, R. F., J. W. Burton, and P. Kwanyuen. (1989). In: Edible Fats and Oils Processing: Basic Principles and Modern Practices. Ed. Erickson, D. AOCS Press, Champaign, pp. 355-359.
- Wu, J. Muir A.D. (2008). Comparative structural, emulsifying, and biological properties of 2 major canola proteins, cruciferin and napin. *J. Food Sci.* **73**: C210–C216.
- Xin-Qing, X., Viet, H. T., Martin, P., Keith, W., Philip, S., (1999). Chemical and physical analyses and sensory evaluation of six deep-frying oils. *J. Am. Oil Chem. Soc.* **76**:1091–1099.

Younis, T. M., Sherazi, S. T. H., Mahesar, S.A., and Ahmed K. A. (2009). Effects of chicken frying on soybean, sunflower and canola oils. *Pak. J. Anal. Environ. Chem.* **10**: 59-66.